

# Convex Team Logics

Alexi Anttila & Søren Brinck Knudstorp

ILLC, University of Amsterdam


Workshop on the Occasion of Marco Degano's Doctoral Defense

# Plan for the talk

- Convexity: What is it and why is it interesting?
- Team Logics: Connectives and notions of propositionhood.
- Results: Expressive completeness for convex team logics.

# Convexity: the why and what

Degano, 2024: *The underlying idea is that the meaning of expressions should denote a **convex** 'region' provided a suitable notion of meaning space. **Convexity would be violated when gaps are present** in the underlying 'region' that expressions denote.*




convex



not convex

Image from Gärdenfors, *The Geometry of Meaning: Semantics Based on Conceptual Spaces*, 2000



# Convexity as Linguistic/Cognitive Universal

## 1. Generalized quantifiers:

Barwise & Cooper, 1981: *The simple NP's of any natural language express monotone quantifiers or **conjunctions of monotone quantifiers**.*

Van Benthem, 1984: *Monotonicity is a strong condition, whose validity for arbitrary logical constants is debatable. Nevertheless, one does expect a certain "smooth" behaviour of reasonable quantifiers; and, therefore, the following notion of **continuity [ed: convexity]** has a certain interest. . .*

## 2. Concept formation:

Gärdenfors, 2000: *A central feature of our cognitive mechanisms is that we assign properties to the objects that we observe [...] I primarily want to pin down the properties that are, in a sense, natural to our way of thinking [...] The third and most powerful criterion of a region is the following, which also relies on betweenness: A subset  $C$  of a conceptual space  $S$  is said to be **convex** if, for all points  $x$  and  $y$  in  $C$ , all points between  $x$  and  $y$  are also in  $C$ .*

# Convexity as Linguistic/Cognitive Universal

## 3. Indefinites:

Degano, 2024: *We can then provide a more grounded explanation for the absence of indefinites that lexicalize only the SK and NS functions as a **violation of the convexity constraint**.*

### Definition (Convexity over Teams)

A set of teams  $\mathcal{P}$  is **convex** iff for all  $t, t', t''$  such that  $t \subseteq t' \subseteq t''$ , if  $t \in \mathcal{P}$  and  $t'' \in \mathcal{P}$ , then  $t' \in \mathcal{P}$ .

## (Propositional) team logics: connectives

Traditionally (in, e.g., CPC), formulas  $\varphi$  are evaluated at **single valuations**  
 $v : \mathbf{Prop} \rightarrow \{0, 1\}$ ,

$$v \models \varphi.$$

In team semantics, formulas  $\varphi$  are evaluated at **sets ('teams') of valuations**  
 $t \subseteq \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$ ,

$$t \models \varphi.$$

### Definition (some team-semantic clauses)

For  $t \subseteq \{v \mid v : \mathbf{Prop} \rightarrow \{0, 1\}\}$ , we define

$t \models p$	<b>iff</b>	$\forall v \in t : v(p) = 1,$
$t \models \varphi \wedge \psi$	<b>iff</b>	$t \models \varphi$ <i>and</i> $t \models \psi,$
$t \models \varphi \vee \psi$	<b>iff</b>	there exist $t', t''$ such that $t' \models \varphi;$ $t'' \models \psi;$ and $t = t' \cup t'',$
$t \models \varphi \wp \psi$	<b>iff</b>	$t \models \varphi$ <i>or</i> $t \models \psi.$

# New connectives

## On connectives:

*Fact 1:* Team semantics for  $\{\neg, \wedge, \vee\}$  gives us **classical logic**.

*Fact 2:* In classical logic,  $\{\neg, \wedge, \vee\}$  is famously **functionally complete**: all other connectives are definable by these.

*Fact 3:* In team semantics,  $\{\neg, \wedge, \vee\}$  can only capture a fraction of the expressible connectives. For example,  $\bowtie$  is not definable using  $\{\neg, \wedge, \vee\}$ .


*Consequence:* We have a semantic framework for expressions beyond classical assertions, such as questions.

**Take-away:** *Teams provide for ways to express meanings not readily expressible in single-valuation semantics; and thus for **considering new connectives!***

# (Propositional) team logics: propositionhood

- Given any condition-based semantics, we obtain a notion of propositionhood defined as a set of conditions. *Slogan:* **Proposition = a set of conditions.**
- In team semantics, conditions are teams.
- So, **propositions are sets of teams.** **Caveat:** The standard terminology is not ‘propositions’ but ‘properties’.

## Example



Since our meaning space now has structure (as powersets), we can consider natural restrictions on what a proposition is. Or what **different kinds of propositions/meanings** there are! For instance, assertions contra questions. (Note the analogy with generalized quantifiers.)



# Notions of propositionhood (closure properties)

**Take-away:** Teams provide for ways to express meanings not readily expressible in single-valuation semantics; and thus for *considering new notions of propositionhood!*

## Definition (some restrictions on propositionhood)

$\phi$ is <i>downward closed</i> :	$[s \models \phi \text{ and } t \subseteq s] \implies t \models \phi$
$\phi$ is <i>union closed</i> :	$[s \models \phi \text{ for all } s \in S \neq \emptyset] \implies \bigcup S \models \phi$
$\phi$ has the <i>empty team property</i> :	$\emptyset \models \phi$
$\phi$ is <i>flat</i> :	$s \models \phi \iff \{v\} \models \phi \text{ for all } v \in s$
$\phi$ is <i>convex</i> :	$[s \models \phi, u \models \phi \text{ and } s \subseteq t \subseteq u] \implies t \models \phi$

Convexity generalizes downward closure:

$$\text{downward closed} \implies \text{convex}$$

# Interface of connectives and propositionhood

The choice of connectives and the corresponding notion of propositionhood are closely connected. Here are some examples:

- Classical formulas are flat (so union closed) [i.e., classical assertions are flat]
- Formulas with  $\sqcup$  might not be union closed. [i.e., questions are not union closed]
- Consider the *epistemic might* operator  $\blacklozenge$ , defined as

$$s \models \blacklozenge \phi \iff \exists t \subseteq s : t \neq \phi \ \& \ t \models \phi.$$

Formulas with  $\blacklozenge$  are not downward closed [i.e., epistemic uncertainty is not persistent]

# Convexity

**Recall Degano, 2024:** *The underlying idea is that **the meaning of expressions** should denote a convex 'region' **provided a suitable notion of meaning space***

**To summarize, we paraphrase:** *The underlying idea is that  $\|\varphi\|$  should denote a convex 'region': **if  $s, u \in \|\varphi\|$  and  $s \subseteq t \subseteq u$ , then  $t \in \|\varphi\|$***

# Expressive completeness

We answer an open question concerning the expressive power of a certain propositional team logic by showing it is capable of capturing the full range of **convex and union-closed** propositions (properties). We also find logics capable of expressing all **convex** propositions.

We say a logic  $L$  is **expressively complete** for a class of properties  $\mathbb{P}$  ( $\|L\| = \mathbb{P}$ ) if

- (i)  $\|L\| \subseteq \mathbb{P}$ : each property  $\|\phi\|$  (where  $\phi \in L$ ) is in  $\mathbb{P}$
- (ii)  $\mathbb{P} \subseteq \|L\|$ : each property  $\mathcal{P} \in \mathbb{P}$  can be expressed by a formula of  $L$ :  $\mathcal{P} = \|\phi\|$  where  $\phi \in L$ .

Example: Propositional dependence logic is expressively complete for the class of **downward-closed (propositional) team properties**

$$\mathbb{D} = \{\mathcal{P} \mid [t \in \mathcal{P} \ \& \ s \subseteq t] \implies s \in \mathcal{P}\}$$

Propositional inquisitive logic is also expressively complete for  $\mathbb{D}$ .

We consider one propositional logic complete for the class of **convex and union-closed (propositional) team properties**

$$\mathbb{CU} = \{\mathcal{P} \mid [[s, u \in \mathcal{P} \ \& \ s \subseteq t \subseteq u] \implies t \in \mathcal{P}] \ \& \ [s, u \in \mathcal{P} \implies s \cup u \in \mathcal{P}]\}.$$

This logic is the propositional fragment of Aloni's **Bilateral State-based Modal Logic**.

We also consider two logics complete for the class of **convex (propositional) team properties**

$$\mathbb{C} = \{\mathcal{P} \mid [s, u \in \mathcal{P} \ \& \ s \subseteq t \subseteq u] \implies t \in \mathcal{P}\}.$$

These logics are (in a sense) convex variants of the downward-closed logics propositional dependence logic and propositional inquisitive logic.

# A Logic for Convex Union-closed Properties

Syntax of **classical propositional logic (with  $\vee$ )**  $\mathbf{PL}_\vee$

$$\alpha ::= p \mid \perp \mid \neg\alpha \mid \alpha \wedge \alpha \mid \alpha \vee \alpha$$



We extend  $\mathbf{PL}_\vee$  with the **nonemptiness atom**  $\mathbf{NE}$ —syntax of  $\mathbf{PL}_\vee(\mathbf{NE})$ :

$$\phi ::= p \mid \perp \mid \neg\alpha \mid \phi \wedge \phi \mid \phi \vee \phi \mid \mathbf{NE}$$



where  $\alpha \in \mathbf{PL}_\vee$ .

$$\begin{aligned} \{v_q\} &\models p \vee q; \\ \{v_q\} &\not\models (p \wedge \mathbf{NE}) \vee (q \wedge \mathbf{NE}) \end{aligned}$$

$$t \models \mathbf{NE} \iff t \neq \emptyset$$

Aloni's (2022) **Bilateral State-based Modal Logic** is a modal extension of  $\mathbf{PL}_\vee(\mathbf{NE})$  (and is similarly complete for convex union-closed modal team properties in the modal setting). Aloni uses  $\mathbf{NE}$  to model a process of pragmatic enrichment which is then used to account for free choice inferences and other phenomena. E.g.,:

You may have coffee or tea  $\rightsquigarrow$  You may have coffee and you may have tea.

$$\diamond((c \wedge \mathbf{NE}) \vee (t \wedge \mathbf{NE})) \models \diamond c \wedge \diamond t$$

To show  $\mathbf{PL}_V(\mathbf{NE}) = \mathbf{CU}$ , we show:

(i)  $\|\mathbf{PL}_V(\mathbf{NE})\| \subseteq \mathbf{CU}$ : by induction.

(ii)  $\mathbf{CU} \subseteq \|\mathbf{PL}_V(\mathbf{NE})\|$ : by constructing **characteristic formulas** for properties in  $\mathbf{CU}$ .

Characteristic formulas for valuations and teams:

$$\chi_v := \bigwedge \{p \mid v \models p\} \wedge \bigwedge \{\neg p \mid v \not\models p\}$$

$$w \models \chi_v \iff w = v$$

$$\chi_s := \bigvee_{v \in s} \chi_v$$

$$t \models \chi_s \iff t \subseteq s$$

Characteristic formulas for flat (downward- and union-closed) properties:

$$t \models \bigvee_{s \in \mathcal{P}} \chi_s \iff t \subseteq \bigcup \mathcal{P}$$

Characteristic formulas for upward-closed properties:

$$t \models \bigwedge_{v_1 \in t_1, \dots, v_n \in t_n} (((\chi_{v_1} \vee \dots \vee \chi_{v_n}) \wedge \mathbf{NE}) \vee \top) \iff \exists s \in \mathcal{P} = \{t_1, \dots, t_n\} : s \subseteq t$$

Characteristic formulas for convex union-closed properties:

$$t \models \bigvee_{v \in s} \chi_v \wedge \bigwedge_{v_1 \in t_1, \dots, v_n \in t_n} (((\chi_{v_1} \vee \dots \vee \chi_{v_n}) \wedge \mathbf{NE}) \vee \top) \iff \exists s \in \mathcal{P} = \{t_1, \dots, t_n\} : s \subseteq t \text{ and } t \subseteq \bigcup \mathcal{P}$$

$$\iff t \in \mathcal{P} \text{ (if } \mathcal{P} \in \mathbf{CU})$$

# Logics for Convex Properties

To get a characteristic formula for all convex properties, we can replace the characteristic formula for **flat properties** with a characteristic formula for **downward-closed properties**.

**Flat (downward- and union-closed) properties:**

$$t \models \phi_{\mathcal{P}}^F \iff t \subseteq \bigcup \mathcal{P}$$

**Upward-closed properties:**

$$t \models \phi_{\mathcal{P}}^U \iff \exists s \in \mathcal{P} : s \subseteq t$$

**Downward-closed properties:**

$$t \models \phi_{\mathcal{P}}^D \iff \exists s \in \mathcal{P} : t \subseteq s$$

**Convex union-closed properties:**

$$\begin{aligned} t \models \phi_{\mathcal{P}}^F \wedge \phi_{\mathcal{P}}^U &\iff \exists s \in \mathcal{P} : s \subseteq t \text{ and } t \subseteq \bigcup \mathcal{P} \\ &\iff t \in \mathcal{P} \text{ (if } \mathcal{P} \in \mathbb{CU}) \end{aligned}$$

**Convex properties:**

$$\begin{aligned} t \models \phi_{\mathcal{P}}^D \wedge \phi_{\mathcal{P}}^U &\iff \exists s_1 \in \mathcal{P} : s_1 \subseteq t \text{ and } \exists s_2 \in \mathcal{P} : t \subseteq s_2 \\ &\iff t \in \mathcal{P} \text{ (if } \mathcal{P} \in \mathbb{C}) \end{aligned}$$



Can we simply extend  $\mathbf{PL}_\vee(\text{NE})$  to get  $\phi_{\mathcal{P}}^D$ ? No. It can be shown that if a logic  $L$  can define  $\|\phi \vee \psi\|$  for all convex  $\phi, \psi$  (notation:  $\mathbb{C} \vee \mathbb{C} \subseteq \|\!|L\|\!$ ), then  $\|\!|L\|\! \not\subseteq \mathbb{C}$  (the logic cannot be convex!)

For instance, let  $\mathcal{P}_1 := \{\{v_1\}, \{v_2, v_3\}\}$  and  $\mathcal{P}_2 := \{\{v_1\}\}$ . Then  $\mathcal{P}_1, \mathcal{P}_2 \in \mathbb{C}$ , so  $\mathcal{P}_1 = \|\phi_1\|$  and  $\mathcal{P}_2 = \|\phi_2\|$  for  $\phi_1, \phi_2 \in L$ . We have  $\|\phi_1 \vee \phi_2\| = \{\{v_1\}, \{v_1, v_2, v_3\}\} \notin \mathbb{C}$ , so if  $\mathbb{C} \vee \mathbb{C} \subseteq \|\!|L\|\!$ , then  $\|\!|L\|\! \not\subseteq \mathbb{C}$ .

We had  $\vee$  in  $\mathbf{PL}_\vee(\text{NE})$ , but  $\mathbf{PL}_\vee(\text{NE})$  can only define  $\phi \vee \psi$  for all convex *and union-closed*  $\phi, \psi$ ; this does not violate convexity.  $\mathbb{C}\mathbb{U} \vee \mathbb{C}\mathbb{U} \subseteq \|\!|L\|\!$  need not imply  $\mathbb{C} \vee \mathbb{C} \subseteq \|\!|L\|\!$ .

We must either (1) modify  $\vee$  to force convexity, or (2) replace  $\vee$  with something else (that still allows us to capture all of classical propositional logic). Recall that [propositional dependence logic](#) and [propositional inquisitive logic](#) are complete for  $\mathbb{D}$  and hence can express  $\phi_{\mathcal{P}}^D$ . We employ strategy (1) to produce a convex extension of propositional dependence logic, and (2) to produce a convex logic similar to propositional inquisitive logic.

# Convex Propositional Dependence Logic

Syntax of **propositional dependence logic**  $\mathbf{PL}_\vee(= (\cdot))$ :

$$\phi ::= p \mid \perp \mid \neg\alpha \mid \phi \wedge \phi \mid \phi \vee \phi \mid = (p_1, \dots, p_n, q)$$

where  $\alpha \in \mathbf{PL}_\vee$ .  $\|\mathbf{PL}_\vee(= (\cdot))\| = \mathbb{D}$ , so  $\|\phi_{\mathcal{P}}^D\| \in \|\mathbf{PL}_\vee(= (\cdot))\|$ .

We modify  $\vee$  to force downward closure, and hence convexity. We also replace  $\text{NE}$  with the epistemic might operator  $\blacklozenge$  to still be able to express  $\phi_{\mathcal{P}}^U$ .

Syntax of **classical propositional logic (with  $\forall$ )**  $\mathbf{PL}_\forall$ :

$$\alpha ::= p \mid \perp \mid \neg\alpha \mid \alpha \wedge \alpha \mid \alpha \forall \alpha$$

Syntax of **convex propositional dependence logic**  $\mathbf{PL}_\forall(= (\cdot), \blacklozenge)$ :

$$\phi ::= p \mid \perp \mid \neg\alpha \mid \phi \wedge \phi \mid \phi \forall \phi \mid = (p_1, \dots, p_n, q) \mid \blacklozenge\phi$$

where  $\alpha \in \mathbf{PL}_\forall$ .

$$t \models \phi \forall \psi \iff \exists s \supseteq t : s = s_1 \cup s_2 \ \& \ s_1 \models \phi \ \& \ s_2 \models \psi$$

$$t \models \blacklozenge\phi \iff \exists s \subseteq t : s \neq \emptyset \ \& \ s \models \phi$$

For downward-closed  $\phi, \psi$ :  $\phi \vee \psi \equiv \phi \forall \psi$ , so  $\|\phi_{\mathcal{P}}^D\| \in \|\mathbf{PL}_\forall(= (\cdot), \blacklozenge)\|$ . We can define  $\chi_t$  using  $\forall$ , and define  $\phi_{\mathcal{P}}^U$  for  $\mathcal{P} = \{t_1, \dots, t_n\}$  by:  $\phi_{\mathcal{P}}^U := \bigwedge_{v_1 \in t_1, \dots, v_n \in t_n} \blacklozenge(\chi_{v_1} \forall \dots \forall \chi_{v_n})$ .

# A Convex Logic Similar to Propositional Inquisitive Logic

Syntax of **classical propositional logic** (with  $\rightarrow$ )  $\mathbf{PL}_{\rightarrow}$ :

$$\alpha ::= p \mid \perp \mid \alpha \wedge \alpha \mid \alpha \rightarrow \alpha$$

Syntax of **propositional inquisitive logic**  $\mathbf{PL}_{\rightarrow}(\vee)$ :

$$\phi ::= p \mid \perp \mid \phi \wedge \phi \mid \phi \rightarrow \phi \mid \phi \vee \phi$$

$$t \models \phi \rightarrow \psi \iff \forall s \subseteq t : s \models \phi \text{ implies } s \models \psi$$

$$t \models \phi \vee \psi \iff t \models \phi \text{ or } t \models \psi$$

Like  $\mathbf{PL}_{\vee}$ ,  $\mathbf{PL}_{\rightarrow}$  is flat, and corresponds to standard classical propositional logic. We define  $\neg_i \phi := \phi \rightarrow \perp$ .  $\phi \vee_i \psi := \neg_i(\neg_i \phi \wedge \neg_i \psi)$ . Using these, we can construct  $\chi_t$  as before.  $\|\mathbf{PL}_{\rightarrow}(\vee)\| = \mathbb{D}$ , and  $\phi_{\mathcal{P}}^D$  is definable as

$$\phi_{\mathcal{P}}^D := \bigvee_{t \in \mathcal{P}} \chi_t$$

We again add the epistemic modality  $\blacklozenge$  to capture  $\phi_{\mathcal{P}}^U$ :

$$\phi_{\mathcal{P}}^U := \bigwedge_{v_1 \in t_1, \dots, v_n \in t_n} \blacklozenge(\chi_{v_1} \vee_i \dots \vee_i \chi_{v_n}) \quad (\mathcal{P} = \{t_1, \dots, t_n\})$$

Problem: with  $\blacklozenge$  and  $\forall$ , the logic is no longer convex. If  $\mathbb{C} \forall \mathbb{C} \subseteq \|\mathbb{L}\|$ , then  $\|\mathbb{L}\| \not\subseteq \mathbb{C}$ . E.g.,  $\blacklozenge p \forall q$  is not convex.

Solution: We can have  $\mathbb{F} \forall \mathbb{F} \subseteq \|\mathbb{L}\|$  (where  $\mathbb{F}$  is the class of flat properties) and hence  $\|\phi_{\mathcal{P}}^D\| = \|\bigvee_{t \in \mathcal{P}} \chi_t\| \in \|\mathbb{L}\|$  without having  $\forall$  in the syntax. In fact,  $\forall$  is already uniformly definable for flat  $\phi, \psi$  using  $\rightarrow$  and  $\blacklozenge$ .

Syntax of  $\mathbf{PL}_{\rightarrow}(\blacklozenge)$ :

$$\phi ::= p \mid \perp \mid \phi \wedge \phi \mid \phi \rightarrow \phi \mid \blacklozenge \phi$$

For any  $\{\alpha_k \mid k \in K\} \subseteq \mathbf{PL}_{\rightarrow}$ ,

$$\bigvee_{k \in K}^- \alpha_k := \bigwedge_{k \in K} ((\bigwedge_{j \in K \setminus \{k\}} \blacklozenge \neg \alpha_j) \rightarrow \alpha_k).$$

Then  $\bigvee_{k \in K}^- \alpha_k \equiv \bigvee_{k \in K} \alpha_k$ . We can define  $\phi_{\mathcal{P}}^U$  as before, and  $\phi_{\mathcal{P}}^D$  as:

$$\phi_{\mathcal{P}}^D := \bigvee_{t \in \mathcal{P}}^- \chi_t$$

# Conclusion

- Importance of convexity.
- Notion of propositionhood in team logics.
- Results:  $\mathbf{PL}_V(\mathbf{NE})$  is expressively complete for convex and union-closed properties. A modal analogue of the result shows that Aloni's BSML is expressively complete for modal convex and union-closed properties.
- Results: Two logics expressively complete for all convex properties. One is similar to propositional dependence logic, the other to propositional inquisitive logic.

Thank you!